



CENTRAL FARM SERVICE

Beef Max

40-20 R-300

MEDICATED

A Supplement For All Classes of Beef Cattle

For Improved Feed Efficiency

ACTIVE DRUG INGREDIENT

Monensin (as Monesin Sulfate... (250mg/lb)..... 300g/ton

GUARANTEED ANALYSIS

Crude Protein, not less than.....	40.0 %
This includes not more than	20.0 %
Equivalent protein from non-protein nitrogen.	
Crude Fat, not less than	2.00 %
Crude Fiber, not more than	6.00 %
Calcium(Ca), not less than	7.00 %
Calcium(Ca), not more than.....	8.00 %
Phosphorus(P), not less than	0.50 %
Salt(NaCl) , not less than.....	4.00 %
Salt(NaCl), not more than.....	5.00 %
Potassium(K),not less than	2.50 %
Vitamin A, not less than	40,000 IU/Lb

INGREDIENTS

Plant Protein Products, Processed Grain By-Products, Animal Protein Products, Grain Products, Roughage Products, Forage Products, Urea, Calcium Carbonate, Dicalcium Phosphate, Monocalcium Phosphate, Salt, Potassium Chloride, Ammonium Sulfate, Vitamin A Supplement, D-Activated Animal Sterol (Source of Vitamin D3) Vitamin E Supplement, Magnesium Oxide, Calcium Sulfate, Manganous Oxide, Ferrous Carbonate, Zinc Oxide, Copper Sulfate, Sodium Selenite, Ethylenediamine Dihydriodide, Cobalt Carbonate.

DIRECTIONS FOR USE

Must be thoroughly mixed with grain and roughage before feeding. Feed at the rate of 1 lb. Per head per day to provide 150 mg of Monensin Sodium. Feed continuously so that each animal consumes not less than 50mg or more than 360 mg of Monensin per head per day.

CAUTION: Do not allow horses & other equines access to feed containing monensin. Ingestion of monensin by equines has been fatal. Monensin-medicated cattle feed is safe for use in cattle only. Do not feed to lactating dairy cows. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. Feeding undiluted or mixing errors resulting in high concentrations of monensin could be fatal to cattle. Must be thoroughly mixed in feeds before use; **do not feed undiluted.**

B 40-20R300

745962

SEB 02/15/16

Manufactured by

Central Farm Service
Truman, MN 56088

Net Wt 50 lb/22.67 kg or Bulk

MEDICATED



CENTRAL FARM SERVICE

Beef Max

BEEF CONCENTRATE

35-0

FOR GROWING AND FINISHING BEEF CATTLE

----- **GUARANTEED ANALYSIS** -----

Crude Protein	not less than	35.00 %
Crude Fat	not less than	1.00 %
Crude Fiber	not more than	15.00 %
Calcium(Ca)	not less than	6.00 %
Calcium(Ca)	not more than	7.00 %
Phosphorus(P)	not less than	0.80 %
Salt(NaCl)	not less than	3.00 %
Salt(NaCl)	not more than	4.00 %
Magnesium (Mg)	not less than	0.50 %
Potassium (K)	not less than	0.70 %
Copper (Cu)	not less than	150 ppm
Selenium (Se)	not less than	3.30 ppm
Zinc (Zn)	not less than	525 ppm
Vitamin A	not less than	32,000 IU/Lb
Vitamin D-3	not less than	8,000 IU/lb
Vitamin E	not less than	40 IU/lb

----- **INGREDIENTS** -----

Plant Protein Products, Calcium Carbonate, Processed Grain By-Products, Salt, Dicalcium Phosphate, Animal Protein Products, Magnesium Oxide, Cobalt Carbonate, Manganese Sulfate, Ethylenediamine Dihydriodide, Zinc Sulfate, Vitamin D3 Supplement, Copper Chloride, Roughage Products, Vitamin E Supplement, Vitamin A Supplement, Sodium Selenite.

RUMINANT MEAT AND BONE MEAL FREE

----- **DIRECTIONS FOR USE** -----

This supplement is designed to be fed to cattle at the rate of 1 to 2 lbs per head per day. This supplement may be top dressed over feed in the bunk or mixed with grain and/or roughage prior to feeding.

Provide sufficient bunk space so all cattle can eat at the same time.

Cattle should have access to fresh, clean water at all time.

----- **CAUTION: Use only as directed.** Consumption of this product by sheep and goats may result in copper toxicity. -----

B 35-0

745338

SEB 02/15/16

Manufactured by
Central Farm Service
Truman, MN 56088

Net Wt 50 lb/22.67 kg or Bulk



CENTRAL FARM SERVICE

Beef/Max

BEEF CONCENTRATE

35-0 B-400

MEDICATED

Type B Medicated Confined Cattle Feed
FOR GROWING AND FINISHING BEEF CATTLE

For improved feed efficiency in cattle fed in confinement for slaughter.

CAUTION: USE ONLY AS DIRECTED

ACTIVE DRUG INGREDIENT

Lasalocid.....400gm/ton

GUARANTEED ANALYSIS

Crude Protein (MIN).....	35.00 %
Crude Fat (MIN).....	1.00 %
Crude Fiber (MAX).....	15.00 %
Calcium (Ca) (MIN).....	6.00 %
Calcium (Ca) (MAX).....	7.00 %
Phosphorus (P) (MIN).....	0.80 %
Salt(NaCl) (MIN).....	3.00 %
Salt(NaCl) (MAX).....	4.00 %
Magnesium (Mg) (MIN).....	0.50 %
Potassium (K) (MIN).....	0.70 %
Copper (Cu) (MIN).....	150 ppm
Selenium (Se) (MIN).....	3.30 ppm
Zinc (Zn) (MIN).....	525 ppm
Vitamin A (MIN).....	32,000 IU/lb
Vitamin D-3 (MIN).....	8,000 IU/lb
Vitamin E (MIN).....	40 IU/lb

INGREDIENTS

Plant Protein Products, Calcium Carbonate, Processed Grain By-Products, Salt, Dicalcium Phosphate, Animal Protein Products, Magnesium Oxide, Cobalt Carbonate, Manganese Sulfate, Ethylenediamine Dihydriodide, Zinc Sulfate, Vitamin D3 Supplement, Copper Chloride, Roughage Products, Vitamin E Supplement, Vitamin A Supplement, Sodium Selenite.

RUMINANT MEAT AND BONE MEAL FREE

DIRECTIONS

Mix 100 pounds of this lasalocid Type B medicated feed into 1900 lbs of non-medicated feed to obtain one ton of Type C medicated feed for cattle containing 20 g/ton lasalocid. Feed the resulting Type C medicated feed continuously at a rate of 20 pounds to provide 200 mg of lasalocid sodium activity per head per day.

CAUTION

Feeding undiluted or mixing errors may result in an excess lasalocid concentration which could be fatal to cattle or sheep. Do not allow equine access to Type A articles or Type B feeds containing lasalocid as ingestion may be fatal. Safety of lasalocid for use in unapproved species has not been established.

WARNING

A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

B 35-0B400

745436

SEB 02/15/16

Manufactured by

Central Farm Service

Truman, MN 56088

Net Wt 50 lb./22.67 kg or Bulk

MEDICATED



CENTRAL FARM SERVICE

Beef Max

HOLSTEIN BEEF FIN RT400

Type B Medicated Feed

For Use in Cattle Feed Only

DO NOT FEED UNDILUTED

FOR FEEDLOT BEEF CATTLE

For improved feed efficiency and reduction of incidence of liver abscesses caused by *Fusobacterium necrophorum* and *Arcanobacterium (Actinomyces) pyogenes* in cattle fed in confinement for slaughter.

CAUTION: Use only as directed.

ACTIVE DRUG INGREDIENT

MONENSIN (as Monesin sodium)400g/ton
TYLOSIN (as Tylosin phosphate)128g/ton

GUARANTEED ANALYSIS

Crude Protein..... min45.0 %
This includes not more than 19.0% equivalent protein from non-protein nitrogen
Crude Fat min1.00 %
Crude Fiber max15.00 %
Calcium(Ca) min7.50 %
Calcium(Ca) max8.60 %
Phosphorus(P)..... min0.50 %
Salt(NaCl)..... min 1.75%
Salt(NaCl), max 2.25%
Potassium(K)..... min 3.50%
Copper min 190 ppm
Selenium (Se) min 4.10 ppm
Zinc (Zn)..... min 650 ppm
Vitamin A min 40,000 IU/Lb
Vitamin D-3..... min 10,000 IU/Lb
Vitamin E min 50 IU/lb

INGREDIENTS

Plant Protein Products, Calcium Carbonate, Forage Products, Processed Grain By-Products, Urea, Potassium Chloride, Salt, Animal Protein Products, Cobalt Carbonate, Manganese Sulfate, Ethylenediamine Dihydriodide, Zinc Sulfate, Vitamin D3 Supplement, Copper Chloride, Roughage Products, Vitamin E Supplement, Vitamin A Supplement, Sodium Selenite.

RUMINANT MEAT AND BONE MEAL FREE

SEE BACK SIDE OF TAG FOR DIRECTIONS FOR USE,
CAUTION AND WARNING STATEMENTS

B HOLRT400

745445

SEB 02/15/16

Manufactured by
Central Farm Service
Truman, MN 56088

Net Wt. 50 lb./22.67 kg or Bulk

MEDICATED



CENTRAL FARM SERVICE

Beef Max
**HOLSTEIN BEEF FIN
RT400**

DIRECTIONS FOR USE

Each pound of this product contains 200 mg Monensin and 64 mg Tylosin. Mix 140 lbs of this product with 1860 lbs grain and/or forage to provide a complete feed that contains 28 g/ton Monensin and 8.96 g/ton Tylosin. Feed at a rate to provide not more than 360 mg Monensin and not less than 60 nor more than 90 mg Tylosin per head per day.

IMPORTANT: Must be thoroughly mixed into feed before use.

CAUTION:

Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by horses has been fatal. Monensin medicated cattle feed is safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gain may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and the amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

WARNING:

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

B HOLRT400

745445

SEB 2/15/16

Manufactured by
Central Farm Service
Truman, MN 56088
Net Wt. 50 lb./22.67 kg or Bulk

MEDICATED



CENTRAL FARM SERVICE

Beef Max

HOLSTEIN BEEF GROWER RT200

Type B Medicated Feed

For Use in Cattle Feed Only.

**DO NOT FEED UNDILUTED
FOR FEEDLOT BEEF CATTLE**

For improved feed efficiency and reduction of incidence of liver abscessed caused by *Fusobacterium necrophorum* and *Arcanobacterium (Actinomyces) pyogenes* in cattle fed in confinement for slaughter.

CAUTION: Use only as directed

ACTIVE DRUG INGREDIENT

Monensin (as Monesin sodium) 200g/ton
Tylosin (as Tylosin phosphate) 90 g/ton

----- **GUARANTEED ANALYSIS** -----

Crude Protein..... min.....45.00 %
This includes not more than 19.0% equivalent protein from non-protein nitrogen.

Crude Fat min1.00 %
Crude Fiber..... max15.0 %
Calcium(Ca) min4.50 %
Calcium(Ca) max5.50 %
Phosphorus(P)..... min0.75 %
Salt(NaCl) min1.75 %
Salt(NaCl) max2.25 %
Potassium(K) min2.75 %
Copper (Cu) min100.00 ppm
Selenium (Se) min2.65 ppm
Zinc (Zn) min350 ppm
Vitamin A min24,000 IU/Lb
Vitamin D-3..... min8,000 IU/Lb
Vitamin E min32 IU/Lb

INGREDIENTS

Processed Grain By-Products, Plant Protein Products, Calcium Carbonate, Animal Protein Products, Urea, Forage Products, Potassium Chloride, Salt, Dicalcium Phosphate, Cobalt Carbonate, Manganese Sulfate, Vitamin D3 Supplement, Ethylenediamine Dihydriodide, Zinc Sulfate, Roughage Products, Copper Chloride, Vitamin E Supplement, Vitamin A Supplement, Sodium Selenite.

Ruminant Meat and Bone Meal Free

DIRECTIONS FOR USE

Each pound of this product contains 100 mg Monesin and 45 mg Tylosin. Mix 200 lbs of this product with 1800 lbs grain and/or forage to provide a complete feed that contains 20 g/ton Monesin and 9.0 g/ton Tylosin. Feed at a rate to provide not more than 360 mg Monesin and not less than 60 nor more than 90 mg Tylosin per head per day.

IMPORTANT: Must be thoroughly mixed into feed before use.

See Back for Caution & Warning statements

B HOLRT200

745443

SEB 02/15/16

Manufactured by

Central Farm Service

Truman, MN 56088

Net Wt. 50 lb./22.67 kg or Bulk

MEDICATED



CENTRAL FARM SERVICE

Beef Max

HOLSTEIN BEEF GROWER RT200

CAUTION:

Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by horses has been fatal. Monensin medicated cattle feed is safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gain may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and the amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

WARNING:

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

B HOLRT200

745443

SEB 02/15/16

Manufactured by
Central Farm Service
Truman, MN 56088

Net Wt. 50 lb./22.67 kg or Bulk

MEDICATED